References:

1. Ahmed O.M., Mohamad M.A. (2009). Synthesis and anti-tumor activities of some new pyridines and pyrazolo [1, 5-a] Pyrimidine. *Eur Med Chem*, **44**, 3519-3523.

2. Ajam M.A., Kumar B.P.R., Suresh B., (2008) Synthesis and biological screening of 5-{[(4,6-disubstituted pyrimidine-2-yl)thio]methyl}-Nphenyl-1,3,4-thiadiazol-2-amines. *Ind J Pharm Sci*, 672-677.

3. Amr AGE, Mohamed A.M., Mohamed S.F., Hafez N.A.A., Hammam A.G., (2006). Anticancer activities of some newly synthesized pyridine, pyrane, and pyrimidine derivatives. *Bioorg Med Chem*, **14**, 5481–5488.

4. Atwal, K. S.; O'Reilly, B. C.; Gougoutas, J. Z.; (1987). Malley, M. F. Synthesis of Substituted 1, 2, 3, 4-Tetrahydro-6-methyl-2-thioxo-5-pyrimidinecarboxylic Acid Esters. *Heterocycles*, **26**, 1189- 1192.

5. Atwal, K. S.; Swanson, B. N.; Unger, S. E.; Floyd, D. M.; Moreland, S.; Hedberg, A.; O'Reilly, B. C. (1991). An expedient protocol of the Biginelli dihydropyrimidine synthesis using carbonyl equivalents. *J.Med. Chem*., **34**(2), 806-811.

6. Becerro M.A., Uriz M.J., & Turon X., (1994). Trends in space occupation by the encrusting sponge Crambe crambe: variation in shape as a function of size and environment. Mar. Biol. **121**. 301-307.

7. Berlinck, R. G. S., (2002), Natural guanidine derivatives. *Nat. Prod. Rep.*, **19**, 617-649.

8. Cho, H.; Ueda, M.; Shima, K.; Mizuno, A.; Hayashimatsu, M.; Ohnaka, Y.;Hamaguchi, M.; Aisaka, K.; Hidaka, T.; Kawai, M.; Takeda, M.; Ishihara, T.;Funahashi, K.; Satoh, F; Morita, M.; Noguchi, T., (1989), *J.Med. Chem*., **32**, 2399-2406.

9. Dardonville, C.; Rozas, I.; Goya, P.; Giron, R.; Goicoechea, C.; Martın, M. I., (2003), acetylsalicylic acid and morphine, respectively, were isolated in 19th century., *Bioorg. Med. Chem.* **11**, 1283.

10. Folkers, K.; Johnson, T. B. (1933) Researches on Pyrimidines. CXXXVI. The Mechanism of Formation of Tetrahydropyrimidines by the Biginelli Reaction. *J. Am. Chem.Soc.* **55**, 3781-3791.

11. Gangjee A. ,Kurup S., Ihnat M.A., Thorpe JE, Shenoy SS. (2010). Synthesis and biological activity of N^4-phenylsubstituted-6-(2,4-dichloro phenylmethyl)-7H-pyrrolo[2,3-d]pyrimidine-2,4-diamines as vascular endothelial growth factor receptor-2 inhibitors and antiangiogenic and antitumor agents. *Bioorg Med Chem*, **18,** 3575–3587.

12. Giovannoni, M. P.; Vergelli, C.; Ghelardini, C.; Galeotti, N.; Bartolini, A.; DalPiaz, V. J., (2003),[(3-chlorophenyl)piperzinylpropyl]pyridazinones and analogues as potent antinociptive agents., *j. Med. Chem*. **46**, 1055.

13. Hurst, E. W.; Hull, R. (1961). Two new synthetic substances active against viruses of the psittacosis-lymphogranulomatrachoma group *J. Med. Pharm. Chem*., **3,** 215.

14. (a) Kashman, Y.; Hirsh, S.; McConnell, O. J.; Ohtani, I.; Kusumi, T.;Kakisawa, H. Ptilomycalin A: (1989). A novel polycyclic guanidine alkaloid of marine origin. *J. Am. Chem. Soc*., **111**, 8925. (b) Jares-Erijman, E. A.; Sakai, R.; Rinehart, K. L. (1991), Kinetics and mechanism of nickel(II) ion retarded Schiff-base hydrolysis of 2,2'-dipyridylmethylideneaniline. *J. Org. Chem*., **56**, 5712.

15. Mohamed N.R., El-Saidi M.M.T., Alia Y.M., Elnagdi M.H. (2007), Utility of 6-amino-2-thiouracil as a precursor for the synthesis of bioactive Pyrimidine derivatives. *Bioorg Med Chem*, **15,** 6227–6235.

16. Overman, L. E.; Rabinowitz, M. H.; Renhowe, P. A., *J. Am. Chem. Soc*.,(1995). **117**, 2675.

17. (a) Patil, A. D.; Kumar, N. V.; Kokke, W. C.; Bean, M. F.; Freyer, A. J.; De Brosse, C.; Mai, S.; Truneh, A.; Faulkner, D. J.; Carte, B.; Breen, A. L.;Hertzberg, R. P. Johnson, R. K.; Westley, J. W.; Potts, B. C. M. .(1995). Novel Alkaloids from the Sponge Batzella sp.: Inhibitors of HIV gp120-Human CD4 Binding *J.Org. Chem*. **60**, 1182. (b) Patil, A. D.; Freyer, A. J.; Taylor, P. B.; Carte, B.; Zuber,G.; Johnson, R. K.; Faulkner, D. J. (1997). Batzelladines F−I, Novel Alkaloids from the Sponge Batzella sp.: Inducers of $p56^{lck}$-CD4 Dissociation. *J. Org. Chem,* **62**, 1814.

18. Patel, R.; Desai, K.; Chikhalia, K., (2003), Many biological, pharmaceutical and therapeutical activities. *J. Ind. Chem. Soc.* **80**, 138.

19. Rovnyak, G. C.; Atwal, K. S.; Hedberg, A.; Kimball, S. D.; Moreland, S.;Gougoutas, J. Z.; O'Reilly, B. C.;Schwartz, J.; Malley, M. F. .(1992), Dihydropyrimidine calcium channel blockers. 4. Basic 3-substituted-4-aryl-1, 4-dihydropyrimidine-5-carboxylic acid esters. Potent antihypertensive agents. *J. Med. Chem*., **35**, 3254-3263

20. Shutalev, A. D.; Kishko, E. A.; Sivova, N. V.; Kuznetsov, A. Yu. (1998). A New Convenient Synthesis of 5-Acyl-1, 2, 3, 4-tetrahydropyrimidine-2- thiones/one.; *Molecules*, ***3,*** 100-106.

21. Sinder, B. B.; Shi, Z. mesoporous .(1993*).* Molecular sieve mcm-41 catalyzed one-pot synthesis Of 3,4-dihydro-2(1h)-pyrimidinones and –thiones under Solvent-free conditions.; *J Org. Chem*., ***5*8**, 3828.

22. Walsh, T. D., (1990), Prevention of opioids side effects. *J. Pain Symptom Manage*. **5**, 362.

23. Kappe, C. O. (1997). A Reexamination of the Mechanism of the Biginelli Dihydropyrimidine Synthesis. Support for an *N*-Acyliminium Ion Intermediate. *J.Org. Chem*. **62**, 7201

24. Kappe C O. (2000). Recent Advances in the Biginelli Dihydropyrimidine Synthesis. New Tricks from an Old Dog. *Acc Chem Res*. **33**,879–888

25 Kappe C O. (2000). Biologically active dihydropyrimidones of the Biginelli-type — a literature survey. *Eur J Med Chem*. **35**:1043–1052